

# Rekonstruktion bei perinealen Defekten

LUKAS BANKAMP, ADRIEN DAIGELER, JOHANNES TOBIAS THIEL  
TÜBINGEN

1. Chan S et al (2010) Use of myocutaneous flaps for perineal closure following abdominoperineal excision of the rectum for adenocarcinoma. *Colorectal Dis* 12: 555–560
2. Pai AA et al (2022) Analysis of clinical and patient-reported outcomes in post-ELAPE perineal reconstruction with IGAP flap – a 5-year review. *JPRAS Open* 34: 10–20
3. Devulapalli C et al (2016) Primary versus flap closure of perineal defects following oncologic resection: A systematic review and meta-analysis. *Plast Reconstr Surg* 137: 1602–1613
4. Mughal M et al (2013) Reconstruction of perineal defects. *Ann R Coll Surg Engl* 95: 539–544
5. Siccardi MB, Bordoni, D (2024) Anatomy, Abdomen and Pelvis, Perineal Body. In: StatPearls. Treasure Island (FL) ineligible companies. Disclosure: Bruno Bordoni declares no relevant financial relationships with ineligible companies.
6. Beier JP et al (2015) Komplikationsvermeidung in der onkologischen Chirurgie der Becken-/Leistenregion. Kombiniert onkochirurgische und plastisch-rekonstruktive Maßnahmen. *Chirurg* 86: 242–250
7. Chang CC et al (2019) Risk factors for delayed perineal wound healing and its impact on prolonged hospital stay after abdominoperineal resection. *World J Surg Oncol* 17: 226
8. Peirce C, Martin S (2016) Management of the Perineal Defect after Abdominoperineal Excision. *Clin Colon Rectal Surg* 29: 160–167
9. Miller DT et al (2020) Novel creation of a noneverted stoma during ileal conduit urinary diversion: Technique and short-term outcomes. *Urology* 146: 260–264
10. Westbom CM, Talbot SG (2019) An algorithmic approach to perineal reconstruction. *Plast Reconstr Surg Glob Open* 7: e2572
11. Das Gupta K et al (2004) Plastisch-rekonstruktive Therapie bei postonkologischen Defekten im Bereich der Beckenregion. *Chirurg* 75: 1135–1142; quiz 1143–1144
12. Stechl NM et al (2011) Mikrochirurgische Beckenbodenrekonstruktion nach Exenteratio pelvis. Verminderte Morbidität und verbesserte Lebensqualität durch ein interdisziplinäres Konzept. *Chirurg* 82: 625–630
13. Mohr Z et al (2014) Interdisziplinäre, primäre Rekonstruktion des Perineums zur Vermeidung von Wundkomplikationen nach abdominoperinealer Rektumexstirpation. *Zentralbl Chir* 139 Suppl 2: e55–62
14. Vogt PM et al (2006) Plastische Rekonstruktion im Rahmen onkologischer Therapieverfahren: Beckenregion. *Zentralbl Chir* 131: 499–505
15. Kieran I et al (2012) Reconstruction of the pelvis and perineum with a free latissimus dorsi myocutaneous flap: a case report. *Ann R Coll Surg Engl* 94: e254–256
16. Duncan A, Byun S, Paletz J (2020) Free Latissimus dorsi flap with femoral artery end-to-side anastomosis for perineal reconstruction following Fournier’s Gangrene. *Plast Surg Case Stud* 6: 2513826X20964133
17. Windhofer C et al (2011) Perineal reconstruction after rectal and anal disease using the local fascio-cutaneous-infragluteal flap: A new and reliable technique. *Surgery* 149: 284–290
18. Thiel JT et al (2023) Feasibility of Perineal Defect Reconstruction with Simplified Fasciocutaneous Inferior Gluteal Artery Perforator (IGAP) Flaps after Tumor Resection of the Lower Rectum: Incidence and Outcome in an Interdisciplinary Approach. *Cancers (Basel)* 15: 3345
19. Allen RJ, Levine JL, Granzow JW (2006) The in-the-crease inferior gluteal artery perforator flap for breast reconstruction. *Plast Reconstr Surg* 118: 333–339
20. Chrelias T et al (2023) Gluteal propeller perforator flaps: A paradigm shift in abdominoperineal amputation reconstruction. *J Clin Med* 12: 4014
21. Korambayil PM, Allalasundaram K, Balakrishnan T (2010) Perforator propeller flaps for sacral and ischial soft tissue reconstruction. *Indian J Plast Surg* 43: 151–157
22. Windhofer C et al (2012) Local fasciocutaneous infragluteal (FCI) flap for vulvar and vaginal reconstruction: a new technique in cancer surgery. *Int J Gynecol Cancer* 22: 132–138
23. Tham NL et al (2010) The pudendal thigh flap for vaginal reconstruction: optimising flap survival. *J Plast Reconstr Aesthet Surg* 63: 826–831
24. Niranjana NS (2006) Perforator flaps for perineal reconstructions. *Semin Plast Surg* 20: 133–144
25. Preetha Rani K et al (2019) Gracilis myocutaneous flap: Adding to the armamentarium of complex sacrococcygeal defect reconstruction. *Indian J Plast Surg* 52: 246–249
26. Daigeler A et al (2011) The versatility of the pedicled vertical rectus abdominis myocutaneous flap in oncologic patients. *Langenbecks Arch Surg* 396: 1271–1279

27. Kim E et al (2022) Abdominal and perineal hernia rates following vertical rectus abdominis myocutaneous (VRAM) flap reconstruction – a supraregional experience. *J Plast Reconstr Aesthet Surg* 75: 1158–1163
28. Vogt PM et al (2001) Die temporäre arterio-venöse Gefäßschlinge (a-v-Loop) als Empfängergefäß für freie Lappentransplantate ohne lokale Anschlußmöglichkeit. In: *Panta Rhei – Umdenken*. Springer, Berlin / Heidelberg
29. Matschke J et al (2019) AV loop free flap: an interdisciplinary approach for perineal and sacral defect reconstruction after radical oncological exenteration and radiation in a colorectal cancer patient. *World J Surg Oncol* 17: 154